

UCRL-JC-133805


PREPRINT

# Human Folate Metabolism Using $^{14}\text{C}$ -Accelerator Mass Spectrometry

A.J. Clifford, A. Arjomand, S.R. Dueker, H. Johnson,  
P.D. Schneider, R.A. Zulim, B.A. Buchholz and J.S. Vogel

This paper was prepared for submittal to the  
*Synthesis and Applications of Isotopes and Isotopically Labeled Compounds 1997*  
*Philadelphia, PA*  
*September 14-18, 1997*

March 25, 1999



This is a preprint of a paper intended for publication in a journal or proceedings.  
Since changes may be made before publication, this preprint is made available with  
the understanding that it will not be cited or reproduced without the permission of the  
author.

#### DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

## HUMAN FOLATE METABOLISM USING $^{14}\text{C}$ -ACCELERATOR MASS SPECTROMETRY

A. J. Clifford<sup>1</sup>, A. Arjomand<sup>1</sup>, S. R. Dueker<sup>1</sup>, H. Johnson<sup>2</sup>, P. D. Schneider<sup>3</sup>, R. A. Zulim<sup>3</sup>, B. A. Buchholz<sup>4</sup>, and J. S. Vogel<sup>4</sup>

Departments of Nutrition<sup>1</sup> and Animal Science<sup>2</sup>, University of California, Davis, CA 95616, Cancer Center<sup>3</sup>, University of California Davis Medical Center, Sacramento, CA 95617, and Center for Accelerator Mass Spectrometry<sup>4</sup>, Lawrence Livermore National Laboratory, Livermore, CA 94551.

Folate is a water soluble vitamin required for optimal health, growth and development. It occurs naturally in various states of oxidation of the pteridine ring and with varying lengths to its glutamate chain. Folates function as one-carbon donors through methyl transferase catalyzed reactions. Low-folate diets, especially by those with suboptimal methyltransferase activity, are associated with increased risk of neural tube birth defects in children, hyperhomocysteinemic heart disease, and cancer in adults. Rapidly dividing (neoplastic) cells have a high folate need for DNA synthesis. Chemical analogs of folate (antifolates) that interfere with folate metabolism are used as therapeutic agents in cancer treatment.

Although much is known about folate chemistry, metabolism of this vitamin *in vivo* in humans is not well understood. Since folate levels in blood and tissues are very low and methods to measure them are inadequate, the few previous studies that have examined folate metabolism used large doses of radiolabeled folic acid in patients with Hodgkin's disease and cancer (Butterworth *et al.* 1969, Krumdieck *et al.* 1978). A subsequent protocol using deuterated folic acid was also insufficiently sensitive to trace a physiologic folate dose (Stites *et al.* 1997). Accelerator mass spectrometry (AMS) is an emerging bioanalytical tool that overcomes the limitations of traditional mass spectrometry and of decay counting of long lived radioisotopes (Vogel *et al.* 1995). AMS can detect attomolar concentrations of  $^{14}\text{C}$  in milligram-sized samples enabling *in vivo* radiotracer studies in healthy humans. We used AMS to study the metabolism of a physiologic 80 nmol oral dose of  $^{14}\text{C}$ -folic acid (1/6 US RDA) by measuring the  $^{14}\text{C}$ -folate levels in serial plasma, urine and feces samples taken over a 150-day period after dosing a healthy adult volunteer.

### MATERIALS AND METHODS

L- $^{14}\text{C}$ (U)-Glutamic acid was purchased from Moravek Biochemicals. Other chemicals were obtained from Sigma Chemical Co. and were reagent grade or better.

Pteroyl- $^{14}\text{C}$ (U)-glutamic acid ( $^{14}\text{C}$ -folic acid) was synthesized according to the method of Plante *et al.* (1980) with some modifications. To 0.789 mmol pteronic acid was added 10 mL trifluoroacetic acid (TFA). Upon dissolution, 10 mL trifluoroacetic anhydride was added and the solution refluxed for 1.5 h. The solvent was evaporated and the oil suspended in 10 mL water by vigorous shaking. The white precipitate was collected on filter paper and dried to yield 0.75 mmol of N<sup>10</sup>-

trifluoroacetylpteronic acid. To make the ethyl ester of the  $^{14}\text{C}$ -glutamic acid, 250  $\mu\text{Ci}$  [ $^{14}\text{C}(\text{U})$ ]-glutamic acid (S.A. 200 mCi/mmol) was diluted with 0.130 mmol glutamic acid and dried. The solid [ $^{14}\text{C}(\text{U})$ ]-glutamic acid was suspended in 3 mL 1% acetyl chloride in ethanol and incubated for 2 h at  $60^\circ\text{C}$ . The solvent was removed to yield [ $^{14}\text{C}(\text{U})$ ]-glutamate diethyl ester-HCl. The diethyl-glutamate was coupled to the  $\text{N}^{10}$ -trifluoroacetylpteronic acid by a mixed anhydride synthesis using isobutylchloroformate in dimethylformamide. The free acid form was restored by hydrolysis in 0.2N NaOH and the pH adjusted to 3.5 with acetic acid. The solution crystallized overnight at  $4^\circ\text{C}$ . The yellow solid was collected by filtration and dried.

The yellow solid was dissolved in TFA and pteroyl-[ $^{14}\text{C}(\text{U})$ ]-glutamic acid (Fig. 1) was isolated by reverse-phase HPLC using a semi-prep column (Ultacarb 5, ODS 30, 250 x 10 mm, Phenomenex, Torrance, CA) and isocratic mobile phase of 850 mL 0.1% TFA in water with 150 mL  $\text{CH}_3\text{CN}$ ; pumped at 1.0 mL/min. Eluant was removed under vacuum and pteroyl-[ $^{14}\text{C}(\text{U})$ ]-glutamic acid stored at  $4^\circ\text{C}$ . Pteroyl-[ $^{14}\text{C}(\text{U})$ ]-L-glutamic co-eluted with standards in reverse-phase and normal phase HPLC systems. The specific activity of the pteroyl-[ $^{14}\text{C}(\text{U})$ ]-L-glutamic acid was 1.25 mCi/mmol.




Fig. 1. Pteroyl-[ $^{14}\text{C}(\text{U})$ ]-L-glutamic acid

Pre-dose urine and feces were collected a day before dosing. Pre-dose blood was drawn just before an 80 nmol dose of the  $^{14}\text{C}$ -folic acid (100 nCi) in ~ 25 mL water was drunk by the fasting volunteer. After dosing, 24 blood samples were drawn day 1 and 18 more over the next 42 d. Urine and feces were collected for 42 days.

Plasma and erythrocytes were separated by centrifugation. Feces were homogenized in 1L 0.5M KOH. Sample aliquots were packaged in a clean room at UC Davis and transported to Lawrence Livermore National Laboratory (LLNL) for AMS. Total carbon was measured with Carlo Erba 1500 NCS analyzer.

Samples were combusted and reduced to graphite (Vogel 1992) for AMS measurement (Vogel and Turteltaub, 1992, Creek *et al.* 1994). Natural  $^{14}\text{C}$  was subtracted from measured isotope ratios. Excess isotope concentration was converted to folate equivalents (parent compound and all metabolites) using the specific activity of the folic acid, its molecular weight and tissue carbon content.

## RESULTS AND DISCUSSION

AMS measured isotope ratios to  $\pm 3\%$  for all samples. Concentration detection limits for plasma, urine and feces were 0.4, 0.04 and 0.12 fmol  $^{14}\text{C}$  folate/mL plasma, based on double the uncertainty in the background. These respective concentrations correspond to absolute detection limits of 0.008, 0.004 and 0.006 fmol  $^{14}\text{C}$  folate in neat plasma, urine and feces.

The  $^{14}\text{C}$  first appeared in plasma 10 min after dosing, peaked at 1-2 h, and was seen in plasma 42 days after dosing (Fig. 2). The asymmetric shape of the plasma  $^{14}\text{C}$ -folate peak suggests a more slowly responding metabolite ( $^{14}\text{C}$ -5-methyltetrahydrofolic acid) may also be present. This compound is being isolated by HPLC and its  $^{14}\text{C}$  will be assayed by AMS.




Fig. 2. Changes in plasma  $^{14}\text{C}$ -folate concentration by time since dosing.

Based on the recovery of 9 nmol  $^{14}\text{C}$ -folate in the first two stools, dose bioavailability was  $\sim 90\%$ . Cumulative urine and feces recoveries were 9 and 11 nmol  $^{14}\text{C}$ -folate, accounting for a quarter of the dose by day 42. At that time urine and feces losses were 0.14 and 0.03 nmol  $^{14}\text{C}$ -folate/d.

## CONCLUSION

High sensitivity of AMS allowed determination of minute quantities of folate not previously measureable in humans. Synthesized pteroyl- $^{14}\text{C}(\text{U})$ -L-glutamic acid had a modest specific activity (1.25 mCi/mmol) that could be followed in blood for six months after a physiologic dose. AMS avoids need for high specific activities and pharmacological doses as needed with scintillation counting. The volunteer received an effective radiation dose of only 1.1 mrem. The low levels of  $^{14}\text{C}$  require

fastidious lab practices to avoid contamination. Generated lab wastes may legally be non-radioactive in many cases. Results will better define folate needs for optimum health.

#### ACKNOWLEDGMENT

Work was performed under the auspices of US Department of Energy at LLNL under contract W-7405-Eng-48. Support was provided by the University of California CLC Program, the LLNL UC-DRD Program, NIH DK 45939 and NIH DK-53081.

#### REFERENCES

Butterworth CE, Baugh CM and Krumdieck C (1969). A Study of Folate Absorption and Metabolism in Man Utilizing Carbon-14-Labeled Polyglutamates Synthesized by Solid Phase Method. *J. Clin. Investig.*, **48**, 1131-1142.

Creek MR, Frantz CE, Fultz E, Haack K, Redwine K, Shen N, Turteltaub KW and Vogel JS (1994). C-14 AMS; Quantification of Biomolecular Interactions Using Microbore and Plate Separations. *Nucl. Instrum. Methods Physics (Research Section B-Beam Interactions with Materials and Atoms)*, **92**, 454-458.

Krumdieck CL, Fukushima K, Fukushima T, Shiota T and Butterworth CE (1978). Long-term Study of the Excretion of Folate and Pterins in a Human Subject after Ingestion of  $^{14}\text{C}$  Folic Acid, with Observations on the Effect of Diphenylhydantoin Administration. *Am. J. Clin. Nutrition*, **31**, 88-93.

Plante LT, Williamson KL and Pastore EJ (1980). Preparation of Folic Acid Specifically Labeled with Carbon-13 in the Benzoyl Carbonyl. *Methods in Enzymology*, **66**, 533-535.

Stites TE, Bailey LB, Scott KC, Toth JP, Fisher WP and Gregory JF III (1997). Kinetic Modeling of Folate Metabolism Through Use of Chronic Administration of Deuterium-Labeled Folic Acid in Men. *Am. J. Clin. Nutrition*, **65**, 53-60.

Vogel JS (1992). Rapid Production of Graphite without Contamination for Biological AMS. *Radiocarbon*, **34**, 344-350.

Vogel JS and Turteltaub KW (1992). Biomolecular Tracing Through Accelerator Mass Spectrometry. *Trends Anal. Chem.*, **11** 142-149.

Vogel JS, Turteltaub KW, Finkel R and Nelson DE (1995). Accelerator Mass Spectrometry - Isotope Quantification at Attomole Sensitivity. *Anal. Chem.*, **67**, A353-A359.